

Peripheral Collisions with STAR

Spencer Klein, Joakim Nystrand and the STAR Collaboration

When heavy nuclei pass each other at large impact parameters, their electromagnetic fields can interact with each other in a two-photon interaction. These reactions produce a variety of final states, from lepton pairs to single mesons to meson and baryon pairs.

Meson coupling to two-photons depends on the meson quark content and isospin; mesons with anomalous coupling may be exotica such as glueballs (states consisting of several gluons), four-quark states ($q\bar{q}q\bar{q}$), hybrids ($q\bar{q}g$) or more radical possibilities. QCD bag model calculations indicate that these states are bound, and there are many experimental candidates[1].

The nuclei can also interact coherently via colorless hadronic interactions, often described as Pomeron exchange; the Pomeron represents the absorptive part of the nuclear potential. Pomerons are also treated as multiple gluon exchange. At RHIC, coherent Pomerons can be observed in the reaction $\gamma P \rightarrow V$ where V is a vector meson. Double-Pomeron interactions may also be seen. Because the Pomerons couple coherently to the entire nucleus, the reaction kinematics are similar to $\gamma\gamma$ interactions[2].

Because the photon flux is proportional to Z^2 , with the Pomeron flux scaling as $A^{4/3}$, the rates for coherent collisions are very high[3]. Below a center of mass energy 1.5 GeV/c, the $\gamma\gamma$ luminosity will be higher than at existing or planned e^+e^- accelerators [3].

STAR can detect these processes by searching for events with 2 or 4 charged particles in the TPC, nothing else visible in the detector (including the forward TPCs), and with total p_\perp less than $\sqrt{2}\hbar c/R$ [4]. The major backgrounds are grazing hadronic nuclear collisions, beam gas events, and photonuclear interactions. Cosmic ray muons and upstream interactions can also appear at the trigger level. Table 1 shows the expected signals and backgrounds for two repre-

sentative analysis channels. The two-track final state $f_2(1270) \rightarrow \pi^+\pi^-$ is a conventional $q\bar{q}$ meson, while the four-track final state $\gamma\gamma \rightarrow \rho^0\rho^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$ is interesting because a bump observed near the $\rho^0\rho^0$ threshold by e^+e^- colliders “has not been explained by models in which only conventional resonances dominate”[1].

Channel	$f_2(1270)$	$\rho^0\rho^0$
Produced	1,900	42
Detected in STAR	660	12
Grazing AA bkgd.	1/5	0.2/0.8
Beam Gas bkgd.	1/1	0.1/0.1
Photonuclear bkgd.	24	1.2
Total Background	26/32	1.5/2.1

Table 1: Signals and backgrounds for gold beams, in 1,000’s of events per year. The numbers before and after the slash are for FRITIOF and VENUS simulations respectively.

References

- [1] R. M. Barnett *et al.* (PDG Group), Phys. Rev. **D54**, 1 (1996), pg. 568.
- [2] S. Klein and E. Scannapieco, “The Gold Flashlight: Coherent Photons (and Pomerons) at RHIC”, LBL-40457, presented at *Photon ’97*, May 10-15, 1997, to be published in the Proceedings.
- [3] S. Klein and E. Scannapieco, STAR Note 243, Feb., 1995 (unpublished).
- [4] J. Nystrand and S. Klein, “Two-Photon Physics at RHIC: Separating Signals from Backgrounds”, LBL-41111, presented at *Hadron ’97*, August 25-30, 1997, to be published in the Proceedings.